


Hough transform



Overview of today's lecture

- Finding boundaries.
- Line fitting.
- Line parameterizations.
- Hough transform.
- Hough circles.
- Some applications.

Slide credits

Most of these slides were adapted from:

- Kris Kitani (15-463, Fall 2016).

Some slides were inspired or taken from:

- Fredo Durand (MIT).
- James Hays (Georgia Tech).

Finding boundaries



Where are the object boundaries?




Human annotated boundaries



Edge detection




Multi-scale edge detection




Edge strength does not necessarily
correspond to our perception of boundaries



Where are the object boundaries?




Human annotated boundaries



Edge detection



Defining boundaries are hard for us too




Where is the boundary of the mountain top?


Lines are hard to find



Original image



Edge detection




Thresholding

Noisy edge image
Incomplete boundaries


Applications




Autonomous Vehicles
(lane line detection)



tissue engineering
(blood vessel counting)



behavioral genetics
(earthworm contours)



Autonomous Vehicles
(semantic scene segmentation)



Computational Photography
(image inpainting)

Line fitting


Line fitting

Given: Many (x_i, y_i) pairs

Find: Parameters (m, c)

Minimize: Average square distance:

$$E = \sum_i \frac{(y_i - mx_i - c)^2}{N}$$



How can we solve this minimization?

Line fitting

Given: Many (x_i, y_i) pairs

Find: Parameters (m, c)

Minimize: Average square distance:


$$E = \sum_i \frac{(y_i - mx_i - c)^2}{N}$$

Using:

$$\frac{\partial E}{\partial m} = 0 \quad \& \quad \frac{\partial E}{\partial c} = 0$$

Note:

$$\bar{y} = \frac{\sum_i y_i}{N} \quad \bar{x} = \frac{\sum_i x_i}{N}$$




$$c = \bar{y} - m \bar{x}$$
$$m = \frac{\sum_i (x_i - \bar{x})(y_i - \bar{y})}{\sum_i (x_i - \bar{x})^2}$$

What are some problems with the approach?

Problems with parameterizations

Where is the line that minimizes E ?


$$E = \sum_{i=1}^n (y_i - mx_i - b)^2$$



Problems with parameterizations

Where is the line that minimizes E ?


$$E = \sum_{i=1}^n (y_i - mx_i - b)^2$$



Problems with parameterizations

Where is the line that minimizes E ?


$$E = \sum_{i=1}^n (y_i - mx_i - b)^2$$



Problems with parameterizations

Where is the line that minimizes E?

$$E = \sum_{i=1}^n (y_i - mx_i - b)^2$$




Line that minimizes E!!


(Error E must be formulated carefully!)

How can we deal with this?

Line fitting is easily setup as a maximum likelihood problem
... but choice of model is important




$$E = \sum_{i=1}^n (y_i - mx_i - b)^2$$




What optimization are we solving here?

Problems with noise



Least-squares error fit



Squared error heavily penalizes outliers

Model fitting is difficult because...



- **Extraneous data:** clutter or multiple models
 - We do not know what is part of the model?
 - Can we pull out models with a few parts from much larger amounts of background clutter?
- **Missing data:** only some parts of model are present
- **Noise**
- **Cost:**
 - It is not feasible to check all combinations of features by fitting a model to each possible subset

So what can we do?

Line parameterizations

Slope intercept form



$$y = mx + b$$


 slope  y-intercept

What are m and b?

Slope intercept form

$$y = mx + b$$

 slope  y-intercept



Double intercept form

$$\frac{x}{a} + \frac{y}{b} = 1$$

x-intercept



y-intercept



What are a and b?

Double intercept form

$$\frac{x}{a} + \frac{y}{b} = 1$$

x-intercept

y-intercept



Derivation:

(Similar slope) $\frac{y - b}{x - 0} = \frac{0 - y}{a - x}$

$$ya + yx - ba + bx = -yx$$

$$ya + bx = ba$$

$$\frac{y}{b} + \frac{x}{a} = 1$$



Normal Form

$$x \cos \theta + y \sin \theta = \rho$$

What are rho and theta?

Normal Form

$$x \cos \theta + y \sin \theta = \rho$$



Derivation:

$$\cos \theta = \frac{\rho}{a} \rightarrow a = \frac{\rho}{\cos \theta}$$

$$\sin \theta = \frac{\rho}{b} \rightarrow b = \frac{\rho}{\sin \theta}$$

plug into: $\frac{x}{a} + \frac{y}{b} = 1$

$$x \cos \theta + y \sin \theta = \rho$$



Hough transform

Hough transform

- Generic framework for detecting a parametric model
- Edges don't have to be connected
- Lines can be occluded
- Key idea: edges **vote** for the possible models

Image and parameter space

variables

$$y = mx + b$$

parameters




Image space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters




Image space

What would a point in image space become in parameter space?

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

a point
becomes a
line



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

two points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

two points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

three points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

three points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

four points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

variables

$$y - mx = b$$

parameters





Image space

four points
become
?



Parameter space

How would you find the best fitting line?





Image space



Parameter space


Is this method robust to measurement noise?

Is this method robust to outliers?

Line Detection by Hough Transform

Algorithm:

1. Quantize Parameter Space (m, c)
2. Create Accumulator Array $A(m, c)$
3. Set $A(m, c) = 0 \quad \forall m, c$
4. For each image edge (x_i, y_i)
For each element in $A(m, c)$
If (m, c) lies on the line: $c = -x_i m + y_i$
Increment $A(m, c) = A(m, c) + 1$
5. Find local maxima in $A(m, c)$



$A(m, c)$

	1					1	
		1				1	
			1		1		
				2			
			1		1		
		1				1	
	1						1

Problems with parameterization

How big does the accumulator need to be for the parameterization (m, c) ?

$A(m, c)$

	1					1		
		1				1		
			1		1			
				2				
			1		1			
		1				1		
	1						1	

The space of m is huge!

$$-\infty \leq m \leq \infty$$

The space of c is huge!

$$-\infty \leq c \leq \infty$$

Better Parameterization

Use normal form:

$$x \cos \theta + y \sin \theta = \rho$$

Given points (x_i, y_i) find (ρ, θ)

Hough Space Sinusoid

$$0 \leq \theta \leq 2\pi$$

$$0 \leq \rho \leq \rho_{\max}$$

(Finite Accumulator Array Size)




Image and parameter space

variables

$$y = mx + b$$

parameters

parameters

$$x \cos \theta + y \sin \theta = \rho$$

variables





Image space



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

parameters

$$x \cos \theta + y \sin \theta = \rho$$

variables




Image space

a point
becomes a
wave



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$






Image space



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$






Image space



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space


variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$

Wait ...why is rho negative?



a line becomes a point




Image space

Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line becomes a point



Parameter space

There are two ways to write the same line:

Positive rho version:

$$x \cos \theta + y \sin \theta = \rho$$

Negative rho version:

$$x \cos(\theta + \pi) + y \sin(\theta + \pi) = -\rho$$

Recall:

$$\sin(\theta) = -\sin(\theta + \pi)$$

$$\cos(\theta) = -\cos(\theta + \pi)$$





Image and parameter space

variables

$$y = mx + b$$

parameters

$$x \cos \theta + y \sin \theta = \rho$$





Image space

a line
becomes a
point



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters





Image space

two points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters





Image space

three points
become
?



Parameter space

Image and parameter space

variables

$$y = mx + b$$

parameters





Image space

four points
become
?




Parameter space

Implementation

1. Initialize accumulator H to all zeros
2. For each edge point (x, y) in the image
For $\theta = 0$ to 180
 $\rho = x \cos \theta + y \sin \theta$
 $H(\theta, \rho) = H(\theta, \rho) + 1$
end
end
3. Find the value(s) of (θ, ρ) where $H(\theta, \rho)$ is a local maximum
4. The detected line in the image is given by
 $\rho = x \cos \theta + y \sin \theta$

H: accumulator array (votes)



NOTE: Watch your coordinates. Image origin is top left!








Image space



Votes

Basic shapes




(in parameter space)



can you guess the shape?

Basic shapes


(in parameter space)




line

Basic shapes


(in parameter space)



line




rectangle




Basic shapes


(in parameter space)



line





rectangle




circle

Basic Shapes



More complex image



In practice, measurements are noisy...





Image space



Votes

Too much noise ...






Image space



Votes


Effects of noise level

Number of votes for a line of 20 points with increasing noise



More noise, fewer votes (in the right bin)

Effect of noise points




More noise, more votes (in the wrong bin)


Real-world example




Original



Edges



parameter space



Hough Lines


Hough Circles

Let's assume radius known

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters


variables

The diagram shows the equation $(x - a)^2 + (y - b)^2 = r^2$. Two red arrows point from the word "parameters" above to the variables a and b . Two green arrows point from the word "variables" below to the variables x and y .

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters

variables

The diagram shows the equation $(x - a)^2 + (y - b)^2 = r^2$. Two red arrows point from the word "parameters" above to the variables a and b . Two green arrows point from the word "variables" below to the variables x and y .

What is the dimension of the parameter space?

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters

variables

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters

variables





Image space



Parameter space

What does a point in image space correspond to in parameter space?

parameters



$$(x - a)^2 + (y - b)^2 = r^2$$

variables

parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$


variables



parameters

$$(x - a)^2 + (y - b)^2 = r^2$$

variables




What if radius is unknown?

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters


variables

The diagram shows the equation $(x - a)^2 + (y - b)^2 = r^2$. Red arrows point from the word "parameters" to the constants a , b , and r . Green arrows point from the word "variables" to the variables x and y .

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters

variables

The diagram shows the equation $(x - a)^2 + (y - b)^2 = r^2$. Red arrows point from the word "parameters" to the constants a , b , and r . Green arrows point from the word "variables" to the variables x and y .

What if radius is unknown?

$$(x - a)^2 + (y - b)^2 = r^2$$

parameters

variables

$$(x - a)^2 + (y - b)^2 = r^2$$


parameters

variables

If radius is not known: 3D Hough Space!

Use Accumulator array $A(a, b, r)$

Surface shape in Hough space is complicated




Using Gradient Information


Gradient information can save lot of computation:

Edge Location (x_i, y_i)

Edge Direction ϕ_i



Assume radius is known:



$$a = x - r \cos \phi$$

$$b = y - r \sin \phi$$

Need to increment only one point in accumulator!

parameters



$$(x - a)^2 + (y - b)^2 = r^2$$

variables

parameters

$$(x - a)^2 + (y - b)^2 = r^2$$

variables



parameters



$$(x - a)^2 + (y - b)^2 = r^2$$


variables

parameters

$$(x - a)^2 + (y - b)^2 = r^2$$

variables






Pennie Hough detector




Quarter Hough detector



Pennie Hough detector



Quarter Hough detector



The Hough transform ...

Deals with occlusion well?



Detects multiple instances?



Robust to noise?



Good computational complexity?



Easy to set parameters?

